

# Finding shortest contractible and shortest separating cycles in embedded graphs\*

Sergio Cabello<sup>†</sup>

February 27, 2009

## Abstract

We give a polynomial-time algorithm to find a shortest contractible cycle (i.e. a closed walk without repeated vertices) in a graph embedded in a surface. This answers a question posed by Hutchinson. In contrast, we show that finding a shortest contractible cycle through a given vertex is NP-hard. We also show that finding a shortest separating cycle in an embedded graph is NP-hard. This answers a question posed by Mohar and Thomassen.

## 1 Introduction

Let  $G$  be a graph embedded in a surface  $\Sigma$ . Closed walks in  $G$  correspond to closed curves in  $\Sigma$ , and hence we may ask for shortest closed walks with certain topological properties, such as being non-contractible. For the rest of this paper the term *cycle* will be used for a *closed walk without repeated vertices*. We need to clarify this because 'cycle' has been used for different concepts in (combinatorial) surfaces [6, 3] and in graph theory [22].

Thomassen [26] (c.f. Mohar and Thomassen [22, Section 4.3]) introduced the *3-path condition*, which can be phrased as follows: a family of cycles  $\mathcal{C}$  satisfies the 3-path condition if the cycles not in  $\mathcal{C}$  generate a subspace of the cycle space that is disjoint from  $\mathcal{C}$ . When  $\mathcal{C}$  satisfies the 3-path condition, then we can find the shortest cycle in  $\mathcal{C}$  in polynomial time using the so-called *fundamental cycle method*. As particular cases, one obtains polynomial-time algorithms to find a shortest non-contractible cycle and a shortest (surface) non-separating cycle in an embedded graph.

The family of contractible cycles does not satisfy the 3-path condition. This naturally leads to the following problem, attributed to Hutchinson in [27] (c.f. [22, Problem 4.3.3.(a)]): is there a polynomial-time algorithm that finds a shortest contractible cycle? Note that the shortest contractible closed walk may use some vertices twice, and hence perhaps it is not a cycle; Figure 1 shows an example in the annulus. The impossibility of visiting the same vertex twice is not a topological concept, and therefore the problem has a combinatorial flavour.

We show that a shortest contractible cycle in an embedded graph can be found in  $O(E^2 \log E)$  time, where  $E$  is the number of edges in the graph. In contrast, we show that it is NP-hard to find the shortest contractible cycle through a given vertex. In the context of graphs on surfaces, it is the first problem for which we cannot find efficiently the shortest

---

\*A preliminary version will be presented at the 20th ACM-SIAM Symposium on Discrete Algorithms (SODA), New York, in January 2009.

<sup>†</sup>Department of Mathematics, IMFM, and Department of Mathematics, FMF, University of Ljubljana, Slovenia. E-mail: [sergio.cabello@fmf.uni-lj.si](mailto:sergio.cabello@fmf.uni-lj.si). Research supported by the Slovenian Research Agency, project J1-7218 and program P1-0297.




Figure 1: A graph embedded in an annulus. A shortest contractible cycle is strictly longer than a shortest contractible closed walk.

object through a given vertex but we can find the shortest object overall. This may have been the main obstacle in previous attempts to tackle the problem. A slight modification of the NP-hardness reduction also shows that it is NP-hard to find a shortest separating cycle. This answers a question by Mohar and Thomassen [22, Problem 4.3.3.(b)].

Our approach consists of using surgery, a standard technique to deal with topological surfaces that cuts the surface along suitable curves. However, surgery introduces copies of vertices, and hence one has to be careful not to use in the resulting surface copies arising from the same vertex. This naturally leads to a closely related problem, which we call *shortest cycle with forbidden pairs*: given a graph  $G$  and a family  $\mathcal{F} \subseteq \binom{V(G)}{2}$  of *forbidden pairs* of vertices, the task is to find a shortest cycle that does not contain any pair of vertices from  $\mathcal{F}$ . This problem is closely related to the shortest path with forbidden pairs, which is NP-hard [14, 18]. Our main contribution is showing that the shortest cycle with forbidden pairs can be solved in polynomial time when  $G$  is a plane graph (a planar graph together with an embedding) and all the vertices in  $\mathcal{F}$  are cofacial. In some sense, the result is tight: the problem becomes NP-hard for plane graphs when all the vertices in  $\mathcal{F}$  can be covered by two faces, instead of just one.

**Related work.** The abstract and the introduction of the paper by Ren and Deng [24] claims to settle the problem of finding a contractible cycle in polynomial time. However, that paper only considers the problem for so-called LEW embeddings, which is a very restricted class of embeddings in which the shortest non-contractible cycle is longer than any facial walk. Closely related is the work by Cabello et al. [3], where the problem of finding a shortest contractible closed walk that encloses an unspecified face is reduced to the problem of finding a minimum cut in an edge-weighted planar graph. This results in a near-linear time algorithm for finding the shortest contractible closed walk.

It turns out that the shortest non-contractible and the shortest non-separating closed walks in an embedded graph are actually cycles, and hence the distinction between cycle and closed walk becomes irrelevant. Thomassen [26] provided the first polynomial-time algorithm to find a shortest non-contractible and a shortest non-separating cycle, running in roughly cubic time. Much work has been done in these problems since then [1, 2, 4, 12, 20]; the current best running time is  $O(\min\{E, g^3\}E \log E)$  for graphs with  $E$  edges embedded in surfaces of genus  $g$ . Chambers et al. [6] considered the problem of finding a shortest splitting closed walk, that is, a non-contractible but separating closed walk that can be infinitesimally perturbed into a simple (injective) curve.

Other shortest ‘objects’ in an embedded graph have also been considered. Erickson and Har-Peled [12] considered the problem of finding a shortest subgraph with the property that

cutting along it leaves a topological disk. Erickson and Whittlesey [13] studied shortest homotopy or homology generators. Colin de Verdière and Lazarus have studied shortest system of loops [8] and pants decompositions [9]. Colin de Verdière and Erickson [7] have studied the problem of finding the shortest walk (closed or not) homotopic to a given one.

**Organization of the paper.** In Section 2 we present some concepts used through the paper. In Section 3 we provide a polynomial-time algorithm to solve the problem of finding a shortest cycle with forbidden pairs in a planar graph when all the vertices in the forbidden pairs are cofacial. In Section 4 we provide a polynomial-time algorithm to find a shortest contractible cycle in an embedded graph. In Section 5 we show NP-hardness of several related problems.

## 2 Background

Our presentation will be using terminology mostly used for graphs embedded in surfaces [22]. An alternative option would be using the terminology of combinatorial surfaces [7].

**Topology.** We review some basic topology of surfaces; see any of the books [16, 21, 25] for a comprehensive treatment. A *surface* (or 2-manifold)  $\Sigma$  is a compact topological space where each point has a neighbourhood homeomorphic to the plane or to a closed halfplane. A point in  $\Sigma$  is a *boundary point* if no neighbourhood of the point is homeomorphic to the plane. The *boundary* of  $\Sigma$  is the union of all boundary points, and it consists of a finite number (possibly 0) of connected components, each component homeomorphic to a circle. A surface is *non-orientable* if it contains a subset homeomorphic to the Möbius band, and *orientable* otherwise. An orientable surface is homeomorphic to a sphere with a number  $g \geq 0$  of handles attached to it and a number  $b \geq 0$  of disjoint open disks removed, for a unique pair  $g, b \geq 0$ . A non-orientable surface is homeomorphic to the connected sum of  $g$  projective planes and a number  $b \geq 0$  of disjoint open disks removed, for a unique pair  $g, b \geq 0$ . In both cases,  $g$  is called the *genus* of the surface and  $b$  is the number of boundary components.

A *path* in  $\Sigma$  is a continuous mapping from  $[0, 1]$  to  $\Sigma$ , a *closed curve* is a continuous mapping from  $\mathbb{S}^1$  to  $\Sigma$ , and an *arc* is a path whose endpoints are on the boundary of  $\Sigma$ . A path, an arc, or a closed curve is *simple* when the mapping is injective.

Two paths or arcs  $p, q$  with  $p(0) = q(0)$  and  $p(1) = q(1)$  are *homotopic* if there is a continuous function  $h : [0, 1]^2 \rightarrow \Sigma$  such that  $p(\cdot) = h(0, \cdot)$ ,  $q(\cdot) = h(1, \cdot)$ ,  $h(\cdot, 0) = p(0)$ , and  $h(\cdot, 1) = p(1)$ . Two closed curves  $\alpha, \beta$  are (freely) *homotopic* if there is a continuous function  $g : [0, 1] \times \mathbb{S}^1 \rightarrow \Sigma$  such that  $\alpha(\cdot) = g(0, \cdot)$  and  $\beta(\cdot) = g(1, \cdot)$ . A closed curve is *contractible* if it is homotopic to a constant mapping.

Cutting the surface along a simple contractible closed curve gives two connected components, and one of them is a topological disk. A closed curve  $\alpha$  is *separating* if cutting the surface along (the image of)  $\alpha$  gives rise to two connected components; the concept is related to  $\mathbb{Z}_2$ -homology. A contractible closed curve is also a separating curve.

Two curves  $\alpha, \beta$  *cross  $c$  times* if the following two conditions hold: (i) there exist infinitesimal continuous perturbations of  $\alpha, \beta$  that cross transversally  $c$  times; and (ii) any infinitesimal continuous perturbations of  $\alpha, \beta$  have at least  $c$  points in common.

**Embeddings.** We assume that the reader is familiar with graphs embedded in surfaces; see [22] for a comprehensive treatment. For surfaces without boundaries, we only consider *cellular embeddings*, where the removal of the image of the graph leaves a set of topological disks. We assume that the embedding is represented in a suitable way, like for example the

gem representation discussed by Eppstein [11]. For our work, it is convenient to treat graphs embedded in surfaces with boundary as those obtained from an embedding in a surface without boundary where the interior of certain faces is removed. Formally, removing the interior of two faces adjacent to a vertex or removing the interior of a face whose facial walk is not a cycle does not give rise to a surface. However, this is a technicality that does not bring problems, and simplifies the exposition. A *plane graph* is a graph embedded in the sphere or in a topological disk.

**Walks and surgery.** A *walk* in a graph  $G$  is a sequence of vertices where two consecutive vertices are connected by an edge in  $G$ . As mentioned before, the term *cycle* refers to a closed walk without repeated vertices. For embedded graphs, we will use the term *arc* for a walk whose end vertices are on the boundary of the surface. For a walk  $\alpha$  through vertices  $x, y$ , we use  $\alpha[x, y]$  to denote the subwalk of  $\alpha$  between  $x, y$ ; if there are multiple appearances of  $x, y$  along  $\alpha$  it will be clear from the context which ones are used. Homotopy of closed walks and cycles is the same as homotopy of closed curves. Homotopy of walks and arcs is the same as homotopy of paths.

Let  $G$  be an embedded graph and let  $\alpha$  be a walk in  $G$ . We use the notation  $G \# \alpha$  to denote the embedded graph obtained after cutting the surface that receives the embedding along the edges of  $\alpha$ . This can be seen as ungluing the two faces adjacent to an edge  $e$  for each edge  $e$  in  $\alpha$ . We denote by  $G \# (\alpha_1, \dots, \alpha_k)$  the embedded graph obtained inductively as  $(G \# (\alpha_1, \dots, \alpha_{k-1})) \# \alpha_k$ .

**Lengths.** We assume that a graph  $G$  has positive edge-weights. The length  $|\alpha|$  of a walk  $\alpha$  is defined as the sum of the weights of the edges along  $\alpha$ , counted with multiplicity. The distance  $d_G(u, v)$  between vertices  $u, v$  is defined as the minimum length over all walks connecting  $u$  to  $v$ . A walk is *tight* if it is shortest in its homotopy class. The following result will be used in our algorithms.

**Lemma 1.** *Let  $G$  be an embedded graph with  $E$  edges.*

- (a) *A shortest non-separating cycle in  $G$  can be found in  $O(E^2 \log E)$  time.*
- (b) *If the surface where  $G$  is embedded has boundary, we can find in  $O(E \log E)$  time a family of tight walks  $\alpha_1, \dots, \alpha_k$  such that  $G \# (\alpha_1, \dots, \alpha_k)$  is a plane graph with  $O(E)$  edges.*

*Proof.* Item (a) is from [12]. Item (b) follows from Lemmas 4 and 5 of [3]; see Theorem 5 of [3].  $\square$

**Shortest cycle with forbidden pairs** Let  $G$  be a graph, not necessarily embedded. A family of forbidden pairs is a collection  $\mathcal{F} \subseteq \binom{V(G)}{2}$ , where  $\binom{A}{2} \subset 2^A$  denotes the collection of subsets of  $A$  with two elements. A walk in  $G$  is *in compliance* with  $\mathcal{F}$  if the following holds: for any forbidden pair  $\{u, v\} \in \mathcal{F}$ , the walk visits at most one of the vertices  $u, v$ . That is, a walk in compliance with  $\mathcal{F}$  cannot visit both vertices in a forbidden pair of  $\mathcal{F}$ . When  $G$  is a plane graph, the family  $\mathcal{F}$  is *cofacial* if all the vertices appearing in  $\mathcal{F}$  are cofacial.

### 3 Shortest cycle with cofacial forbidden pairs

In this section we consider the problem of finding a shortest cycle in a plane graph  $G$  with forbidden pairs  $\mathcal{F}$ , when the vertices in  $\mathcal{F}$  are cofacial. We will assume that  $G$  is 2-connected, as otherwise we can decompose the graph into maximal 2-connected subgraphs (usually called

blocks) in linear time, and treat each block separately. In a 2-connected plane graph all facial walks are cycles.

Let us mention that, given a planar graph, without embedding, and a family of forbidden pairs  $\mathcal{F}$ , we can decide if there is an embedding of  $G$  where  $\mathcal{F}$  is cofacial by considering an augmented graph  $\hat{G}$  obtained from  $G$  by adding a new vertex  $v_{\text{new}}$  with edges to each vertex appearing in  $\mathcal{F}$ . The graph  $\hat{G}$  is planar if and only if  $G$  has an embedding where the vertices of  $\mathcal{F}$  are cofacial. Moreover, the removal of  $v_{\text{new}}$  in any planar embedding of  $\hat{G}$  provides an embedding of  $G$  where all vertices in  $\mathcal{F}$  are cofacial. Note that deciding if a given graph is planar and constructing a planar embedding of a given planar graph can be done in linear time; see for example [19, 23]. Therefore, this procedure takes linear time.

Let  $f_0$  be the face adjacent to all vertices in  $\mathcal{F}$ , and let  $C_0$  be the facial cycle defining  $f_0$ . We may assume that  $G$  is embedded in a disk such that  $C_0$  defines the boundary of the disk. Let  $E(C_0)$  and  $V(C_0)$  denote the edges and vertices of  $C_0$ , respectively.

We next describe a recursive algorithm. The base case is when  $G = C_0$  is a cycle, and we have two options: if  $\mathcal{F}$  is empty, then  $C_0$  is the shortest cycle in compliance with  $\mathcal{F}$ ; if  $\mathcal{F}$  is nonempty, then  $G$  does not contain any cycle in compliance with  $\mathcal{F}$ . For the general case, we would like to cut along any tight arc connecting two arbitrary vertices of  $C_0$ , and obtain two smaller problems. However, it may be that any tight arc between any two vertices of  $C_0$  is contained in  $C_0$ , and hence we cannot get smaller subproblems in such way. We fix this by considering the graph  $H = G - E(C_0)$ , as follows. First note that some pair of distinct vertices of  $V(C_0)$  are connected through a path in  $H$  because  $G$  is 2-connected and  $G$  is not a cycle. Take the distance  $d_H(u, v)$  in  $H$  for each pair of vertices  $u, v \in V(C_0)$ ,  $u \neq v$ , and let  $u_0, v_0$  be a pair minimizing it; that is

$$(u_0, v_0) = \arg \min_{(u,v) \in (V(C_0))^2, u \neq v} d_H(u, v).$$

Let  $\pi_0$  be a shortest path in  $H$  between  $u_0$  and  $v_0$ . Note that  $\pi_0$  is disjoint from  $E(C_0)$  by the definition of  $H$ . We have the following observation.

**Lemma 2.** *Assume that  $G$  contains some cycle in compliance with  $\mathcal{F}$  and that  $C_0$  is not a shortest cycle in  $G$  in compliance with  $\mathcal{F}$ . Then there exists a shortest cycle in  $G$  in compliance with  $\mathcal{F}$  that does not cross  $\pi_0$ .*

*Proof.* If  $\pi_0$  is a shortest path in  $G$ , then the result would be easier; see Lemma 5 below. However,  $\pi_0$  is a shortest path only in  $H$ , which complicates things and leads us to a case analysis. Through the proof we will assume that all cycles have the same orientation, which is clockwise in our figures.

Let  $\mathcal{Q}$  be the class of cycles of  $G$  in compliance with  $\mathcal{F}$  that are shortest. We have  $\mathcal{Q} \neq \emptyset$  because by hypothesis there exist cycles in compliance with  $\mathcal{F}$ . Each cycle of  $\mathcal{Q}$  bounds a topological disk. Let  $C_{\text{short}}$  be a cycle from  $\mathcal{Q}$  that bounds a disk  $Disk_{\text{short}}$  whose closure does not contain any other cycle from  $\mathcal{Q}$ . We will show that  $\pi_0$  is disjoint from the interior of  $Disk_{\text{short}}$ , and therefore  $C_{\text{short}}$  does not cross  $\pi_0$ .

Let us assume, for the sake of contradiction, that  $\pi_0$  is not disjoint from the interior of  $Disk_{\text{short}}$ . Consider a maximal subpath  $\pi_0[x, y]$  with the property that  $\pi_0[x, y] \setminus \{x, y\}$  is contained in the interior of  $Disk_{\text{short}}$ . Note that the endpoints  $x, y$  are vertices of  $C_{\text{short}}$ . Let  $Q_1$  be the closed walk obtained by concatenating  $\pi_0[x, y]$  with  $C_{\text{short}}[y, x]$  and let  $Q_2$  be the closed walk obtained by concatenating  $\pi_0[y, x]$  with  $C_{\text{short}}[x, y]$ . Note that  $Q_1, Q_2$  are cycles, and they are contained in  $Disk_{\text{short}}$ . Moreover,  $Q_1, Q_2$  are in compliance with  $\mathcal{F}$  because  $\pi_0$  does not contain edges from  $C_0$ . We will show that  $Q_1$  or  $Q_2$  is no longer than  $C_{\text{short}}$ , leading to a contradiction with the choice of  $C_{\text{short}}$ . We distinguish cases depending on the overlap of  $C_{\text{short}}$  and  $C_0$ . Note that  $C_{\text{short}} \neq C_0$  by hypothesis, so we have the following three scenarios.

- (i) If  $C_{\text{short}}$  and  $C_0$  do not have any common edge, then  $C_{\text{short}}$  is a shortest cycle in  $H$ . See Figure 2, left. In this case  $|\pi_0[x, y]| \leq |C_{\text{short}}[x, y]|$  because  $\pi_0$  is a shortest path in  $H$ . Therefore

$$|Q_1| = |\pi_0[x, y]| + |C_{\text{short}}[y, x]| \leq |C_{\text{short}}[x, y]| + |C_{\text{short}}[y, x]| = |C_{\text{short}}|,$$

and  $Q_1$  is no longer than  $C_{\text{short}}$ .

- (ii) If  $C_{\text{short}}$  and  $C_0$  agree in a single maximal subpath  $C_{\text{short}}[v_1, u_1]$  of  $C_{\text{short}}$ , then  $v_1, u_1 \in V(C_0)$  and  $C_{\text{short}}[u_1, v_1]$  is contained in  $H$ . We have three subcases:

- $x, y$  are vertices along  $C_{\text{short}}[u_1, v_1]$ . See Figure 2, center. Renaming the vertices  $x, y$  if necessary, we may assume that the ordering of vertices along  $C_{\text{short}}[u_1, v_1]$  is  $u_1, x, y, v_1$ , possibly with  $u_1 = x$  or  $y = v_1$ . Since  $C_{\text{short}}[u_1, v_1]$  is contained in  $H$ , also  $C_{\text{short}}[x, y]$  is contained in  $H$ . We then have  $|\pi_0[x, y]| \leq |C_{\text{short}}[x, y]|$  because  $\pi_0$  is a shortest path in  $H$ , and therefore  $Q_1$  is no longer than  $C_{\text{short}}$ .
- $x, y$  are vertices along  $C_{\text{short}}[v_1, u_1]$ . See Figure 2, right. In this case  $\{x, y\} = \{u_0, v_0\}$ . Renaming the vertices  $x, y$  if necessary, we may assume that the ordering of vertices is such that  $C_{\text{short}}[u_1, v_1] \subseteq C_{\text{short}}[x, y]$ . By the choice of  $u_0, v_0$  we have

$$|\pi_0[x, y]| = |\pi_0[u_0, v_0]| \leq |C_{\text{short}}[u_1, v_1]| \leq |C_{\text{short}}[x, y]|,$$

and therefore  $Q_1$  is no longer than  $C_{\text{short}}$ .

- $x, y$  are not both along  $C_{\text{short}}[u_1, v_1]$  or both along  $C_{\text{short}}[v_1, u_1]$ . See Figure 3, left and center. In this case,  $\{x, y\} \cap \{u_1, v_1\} = \emptyset$ . Renaming the vertices  $u_0, v_0$  if necessary, we may assume that  $v_0$  is along the path  $C_{\text{short}}[v_1, u_1]$ . Renaming the vertices  $x, y$  if necessary, we may assume that  $y = v_0$ , and hence  $x$  is along the path  $C_{\text{short}}[u_1, v_1]$ . When  $u_1 \neq u_0$  (Figure 3, left), the concatenation of  $C_{\text{short}}[u_1, x]$  and  $\pi_0[x, u_0]$  is in  $H$  and connects two different vertices of  $V(C_0)$ . Because of the choice of  $u_0, v_0$  we have

$$|\pi_0[v_0, x]| + |\pi_0[x, u_0]| = |\pi_0[v_0, u_0]| \leq |C_{\text{short}}[u_1, x]| + |\pi_0[x, u_0]|,$$

and hence

$$|\pi_0[y, x]| = |\pi_0[v_0, x]| \leq |C_{\text{short}}[u_1, x]| \leq |C_{\text{short}}[y, x]|.$$

This implies that  $Q_2$  is no longer than  $C_{\text{short}}$ . When  $u_1 = u_0$  (Figure 3, center), then  $v_1 \neq u_0$ , and the same argument used with the concatenation of  $\pi_0[v_0, x]$  and  $C_{\text{short}}[x, v_1]$  shows that  $|\pi_0[x, y]| \leq |C_{\text{short}}[x, y]|$ , and hence  $Q_1$  is no longer than  $C_{\text{short}}$ .

- (iii) If  $C_{\text{short}}$  and  $C_0$  agree in at least two (disjoint) maximal subpaths of  $C_{\text{short}}$ , then there exist vertices  $u_1, v_1, u_2, v_2 \in V(C_0)$  such that the paths  $C_{\text{short}}[u_1, v_1], C_{\text{short}}[u_2, v_2]$  are disjoint from  $C_0$ , except at its endpoints. See Figure 3, right. Since the paths  $C_{\text{short}}[u_1, v_1], C_{\text{short}}[u_2, v_2]$  are contained in  $H$ , the choice of  $u_0, v_0$  implies

$$|C_{\text{short}}[u_1, v_1]| \geq |\pi[u_0, v_0]| \quad \text{and} \quad |C_{\text{short}}[u_2, v_2]| \geq |\pi[u_0, v_0]|.$$

Since the paths  $C_{\text{short}}[u_1, v_1], C_{\text{short}}[u_2, v_2]$  are edge-disjoint, we have

$$|C_{\text{short}}| \geq |C_{\text{short}}[u_1, v_1]| + |C_{\text{short}}[u_2, v_2]| \geq 2 \cdot |\pi[u_0, v_0]|,$$

and hence

$$|C_{\text{short}}[x, y]| + |C_{\text{short}}[y, x]| = |C_{\text{short}}| \geq 2 \cdot |\pi[u_0, v_0]|.$$




Figure 2: Left: an example for case (i) of Lemma 2. Center: an example for the first item in case (ii) of Lemma 2. Right: an example for the second item in case (ii) of Lemma 2.




Figure 3: Left and center: examples for the third item in case (ii) of Lemma 2. Right: an example for case (iii) of Lemma 2.

This means that

$$\max\{|C_{\text{short}}[x, y]|, |C_{\text{short}}[y, x]|\} \geq |\pi[u_0, v_0]| \geq |\pi[x, y]|,$$

and  $Q_1$  or  $Q_2$  is no longer than  $C_{\text{short}}$ .

□

**Theorem 3.** *Given a plane graph  $G$  with  $n$  vertices and a family  $\mathcal{F}$  of  $F$  forbidden pairs whose vertices are all adjacent to one face of  $G$ , we can find a shortest cycle in compliance with  $\mathcal{F}$  in  $O(n^2 + nF)$  time.*

*Proof.* Like before, we assume that  $G$  is 2-connected, that  $C_0$  is the facial cycle of  $G$  containing all vertices in  $\mathcal{F}$ , and that  $G$  is embedded in a disk whose boundary is defined by  $C_0$ . Consider the following recursive algorithm:

- (i) If  $G$  is a cycle and  $\mathcal{F}$  is empty, we return the cycle  $G$ .
- (ii) If  $G$  is a cycle and  $\mathcal{F}$  is nonempty, we return that  $G$  has no cycle in compliance with  $\mathcal{F}$ .
- (iii) If  $G$  is not a cycle, we construct  $H = G - E(C_0)$ , find a shortest path  $\pi_0$  between vertices

$$(u_0, v_0) = \arg \min_{(u,v) \in (V(C_0))^2, u \neq v} d_H(u, v),$$

and cut  $G$  along  $\pi_0$  to obtain two 2-connected plane graphs, denoted  $G_1, G_2$ ; see Figure 4. We regard  $G_1, G_2$  as embedded in topological disks. We also construct two families of forbidden sets  $\mathcal{F}_1, \mathcal{F}_2$  from  $\mathcal{F}$  in the natural way:  $\mathcal{F}_i$  contains each forbidden pair of  $\mathcal{F}$  whose two vertices (or copies of them) are in  $G_i$ . If  $\{u_0, v_0\} \in \mathcal{F}$ , then




Figure 4: Left: the path  $\pi_0$ . Right: the graphs  $G_1, G_2$  obtained after cutting  $G$  along the arc  $\pi_0$ .

$\{u_0, v_0\}$  goes to both  $\mathcal{F}_1, \mathcal{F}_2$ . All other forbidden pairs of  $\mathcal{F}$  go only to  $\mathcal{F}_1$  or to  $\mathcal{F}_2$ . For  $i = 1, 2$ , we *recursively* find a shortest cycle  $C_i$  in  $G_i$  in compliance with  $\mathcal{F}_i$ , or find out that no cycle of  $G_i$  is in compliance with  $\mathcal{F}_i$ . If  $\mathcal{F}$  is empty, we return the shortest among  $C_0, C_1, C_2$ , or the subset of them that is defined. If  $\mathcal{F}$  is nonempty and for  $i = 1, 2$  no cycle of  $G_i$  is in compliance with  $\mathcal{F}_i$ , we then return that no cycle of  $G$  is in compliance with  $\mathcal{F}$ . Otherwise, we return the shortest cycle among  $C_1, C_2$ , or the subset of them that is defined.

Correctness of the algorithm is shown by induction on  $|E(H)|$ . If  $|E(H)| = 0$ , then  $G$  is a cycle, and correctness follows from cases (i) and (ii) of the algorithm. If  $|E(H)| \geq 1$ , then  $G$  is not a cycle, and  $G_1$  and  $G_2$  are also 2-connected plane graphs. For  $i = 1, 2$ , let  $H_i$  be the subgraph of  $G_i$  containing the edges not in the boundary of the disk holding the embedding of  $G_i$ . It clearly holds  $|E(H_i)| < |E(H)|$ , and by the induction hypothesis the algorithm correctly finds a shortest cycle  $C_i$  in  $G_i$  that is in compliance with  $\mathcal{F}_i$ , and hence with  $\mathcal{F}$ , or tells that no such cycle exists. If  $C_0$  is a shortest cycle in compliance with  $\mathcal{F}$ , then the algorithm returns  $C_0$ , and hence the algorithm is correct. Otherwise,  $C_0$  is not a shortest cycle in compliance with  $\mathcal{F}$ , and because of Lemma 2, the shortest cycle between  $C_1, C_2$  is the shortest cycle in compliance with  $\mathcal{F}$ . This finishes the proof of correctness.

Let us now bound the time complexity of the algorithm. Finding the path  $\pi_0$  can be done in linear time, as follows. Firstly, construct the graph  $H = G - E(C_0)$  and add an extra vertex  $v_{\text{new}}$  with edges of length zero to all vertices  $V(C_0)$ . Let  $H'$  be the resulting planar graph. Secondly, construct in  $H'$  a shortest path tree  $T_{sp}$  from the source  $v_{\text{new}}$  in linear time [17]. Each edge  $e \in E(H) \setminus E(T_{sp})$  defines a cycle, denoted by  $\text{cycle}(T_{sp}, e)$ , that is the concatenation of  $e$  with the subpath of  $T_{sp}$  between the endpoints of  $e$ . Let  $e_0$  denote an edge containing the middle point of the path  $\pi_0$ . On the one hand,  $\text{cycle}(T_{sp}, e_0)$  has the same length as  $\pi_0$ . On the other hand, for any cycle  $\text{cycle}(T_{sp}, e)$  that passes through  $v_{\text{new}}$  the path  $\text{cycle}(T_{sp}, e) - v_{\text{new}}$  is a shortest path in  $H$  between two vertices of  $V(C_0)$ . Therefore, we can compute a best edge  $e^*$  as

$$e^* = \arg \min_{e \in E(H) \setminus E(T_{sp}) \text{ s.t. } v_{\text{new}} \in \text{cycle}(T_{sp}, e)} \{|\text{cycle}(T_{sp}, e)|\},$$

and then take  $u_0, v_0$  as the two vertices of  $V(C_0)$  in  $\text{cycle}(T_{sp}, e^*)$ . Finally, we compute the shortest path  $\pi_0$  in  $H$  between  $u_0$  and  $v_0$  in linear time.

Consider a rooted binary tree where each node corresponds to a subproblem considered through the recursive algorithm. The root corresponds to the original problem, and the two children of a node contain disjoint subsets of the faces from its parent. This tree has at most one leaf-node per face of  $G$ , and therefore a total of  $O(n)$  subproblems are considered. In each subproblem we spend  $O(n)$  time to find the path  $\pi_0$  and  $O(F + n)$  time to check if the boundary cycle  $C_0$  is in compliance with  $\mathcal{F}$ . The running time follows.  $\square$




Figure 5: Figure for the proof of Lemma 5.

In our forthcoming application, the forbidden pairs  $\mathcal{F}$  have some additional structure. A partition  $\mathcal{V} = \{V_1, V_2, \dots\}$  of  $V(G)$  defines the following set of forbidden pairs

$$\mathcal{F}(\mathcal{V}) = \bigcup_{V_i \in \mathcal{V}} \binom{V_i}{2}.$$

Here we assume that  $\binom{V_i}{2} = \emptyset$  whenever  $V_i$  has one element. Let us say that a partition  $\mathcal{V}$  is cofacial when  $\mathcal{F}(\mathcal{V})$  is cofacial.

**Corollary 4.** *Given a plane graph  $G$  with  $n$  vertices and a cofacial partition  $\mathcal{V}$  of  $V(G)$ , we can find a shortest cycle in compliance with  $\mathcal{F}(\mathcal{V})$  in  $O(n^2)$  time.*

*Proof.* Given the partition  $\mathcal{V}$  of  $V(G)$ , we can check in  $O(n)$  time if a cycle is in compliance with  $\mathcal{F}(\mathcal{V})$  or not. Hence, in each of the  $O(n)$  subproblems of the recursive algorithm of Theorem 3 we spend  $O(n)$  time, instead of  $O(n + F)$ . The result follows.  $\square$

## 4 Shortest contractible cycle in embedded graphs

Let  $G$  be a graph embedded in a surface, possibly with boundary, and let  $\mathcal{F}$  be a family of forbidden pairs. Here we state the basic tool; see [3, Lemma 3] for a similar statement. The key observation is to consider the case when the vertices appearing in  $\mathcal{F}$  are on the boundary of the surface.

**Lemma 5.** *Let  $\alpha$  be a tight arc or a tight closed walk in  $G$ . If all the vertices of  $\mathcal{F}$  are on the boundary of the surface then there exists a shortest contractible cycle in compliance with  $\mathcal{F}$  that does not cross  $\alpha$ .*

*Proof.* Let  $\mathcal{Q}$  be the class of shortest contractible cycles of  $G$  in compliance with  $\mathcal{F}$ . Each cycle of  $\mathcal{Q}$  bounds a topological disk. Let  $C_{\text{short}}$  be a cycle from  $\mathcal{Q}$  bounding a disk  $Disk_{\text{short}}$  whose closure does not contain any other cycle from  $\mathcal{Q}$ . We will show that  $\alpha$  is disjoint from the interior of  $Disk_{\text{short}}$ , and therefore  $C_{\text{short}}$  does not cross  $\alpha$ .

Assume for the sake of contradiction, that  $\alpha$  enters the interior of  $Disk_{\text{short}}$ ; see Figure 5. Consider a maximal subpath  $\alpha[x, y]$  with the property that  $\alpha[x, y] \setminus \{x, y\}$  is contained in the interior of  $Disk_{\text{short}}$ . Note that the endpoints  $x, y$  are vertices of  $C_{\text{short}}$ . It must be  $x \neq y$ , as otherwise  $\alpha[x, y]$  would be contractible because it would be contained in the topological disk  $Disk_{\text{short}}$ . Similarly,  $\alpha[x, y]$  cannot have any repeated vertices, because the subpath between two repeated vertices would be contractible. It results that the closed walk  $C'$ , defined as the concatenation of  $\alpha[x, y]$  and  $C_{\text{short}}[y, x]$  has no repeated vertices, and hence is a cycle.

Note that the interior of  $Disk_{\text{short}}$  cannot contain any vertex appearing in  $\mathcal{F}$  because  $Disk_{\text{short}}$  does not contain any boundary component. Therefore, the cycle  $C'$  is also in *compliance* with  $\mathcal{F}$  because the ‘new’ subpath  $\alpha[x, y] \setminus \{x, y\}$  does not contain vertices appearing in  $\mathcal{F}$ . Moreover, the paths  $\alpha[x, y]$  and  $C_{\text{short}}[y, x]$  are homotopic because  $\alpha[x, y]$  is inside the disk  $Disk_{\text{short}}$ , and hence  $C'$  is also *contractible*. Using that  $|\alpha[x, y]| \leq |C_{\text{short}}[y, x]|$  because  $\alpha$  is tight, we see also that in fact  $C'$  is *no longer* than  $C_{\text{short}}$ . We conclude that  $C' \in \mathcal{Q}$ . Moreover,  $C'$  is *contained* in the closure of  $Disk_{\text{short}}$  by construction. Therefore the properties of  $C'$  contradict the choice of  $C_{\text{short}}$ .  $\square$

**Theorem 6.** *Given a graph  $G$  with  $E$  edges embedded in a surface, possibly with boundary, we can find in  $O(E^2 \log E)$  time a shortest contractible cycle in  $G$ .*

*Proof.* We first give an algorithm, then discuss its correctness, and finally derive its running time. If the surface has genus 0, we take  $G' = G$ . Otherwise, we compute a shortest non-separating cycle  $\alpha$  in  $G$ , and construct the embedded graph  $G' = G \setminus \alpha$ . Note that  $G'$  has at least one boundary component. We then construct the embedded graph  $G'' = G' \setminus (\alpha_1, \alpha_2, \dots, \alpha_k)$ , where  $\alpha_1, \dots, \alpha_k$  is the tight system of arcs from Lemma 1(b). Note that  $G''$  is a plane graph. We then group the vertices in the boundary of  $G''$  as follows: for each vertex  $v \in V(G)$ , let  $W_v$  be the vertices in  $G''$  that arose as copies of the original vertex  $v$ . Therefore, the groups  $W_v$ ,  $v \in V(G)$ , form a partition of  $V(G'')$ . Note that the only groups in  $W_v$ ,  $v \in V(G)$ , with cardinality larger than one have vertices in the boundary of  $G''$  because copies of vertices are only introduced along the boundary. Therefore,  $\mathcal{F}'' = \mathcal{F}(\{W_v \mid v \in V(G)\}) = \bigcup_{v \in V(G)} \binom{W_v}{2}$  is a cofacial family of forbidden pairs, and we can find the shortest cycle  $C_{\text{short}}$  in  $G''$  in compliance with  $\mathcal{F}''$  using Corollary 3. Finally, we return  $C_{\text{short}}$ . This finishes the description of the algorithm.

We next show the correctness of the algorithm. The embedded graph  $G''$  is obtained iteratively from  $G$  by cutting along a tight walk, until the plane graph  $G''$  is obtained. Let  $H_0, H_1, H_2, \dots, H_t$ , where  $H_0 = G$  and  $H_t = G''$ , be the sequence of embedded graphs constructed through the algorithm, where  $H_{i+1}$  is obtained from  $H_i$  by cutting along one tight walk. For any embedded graph  $H_i$ , let  $\mathcal{F}_i$  be the family of forbidden pairs given by pairs  $\{u, v\} \in \binom{V(H_i)}{2}$  where  $u, v$  are copies of the same vertex in the original graph  $G$ . Note that by construction all the vertices in  $\mathcal{F}_i$  are in the boundary of  $H_i$ . A shortest contractible cycle in compliance with  $\mathcal{F}_{i+1}$  in  $H_{i+1}$  is a shortest contractible cycle in compliance with  $\mathcal{F}_i$  in  $H_i$  because of Lemma 5. By induction, this means that a shortest contractible cycle in compliance with  $\mathcal{F}_i$  in  $H_i$  is a shortest contractible cycle in  $H_0 = G$ . In particular, the cycle  $C_{\text{short}}$ , which is the shortest contractible cycle in compliance with  $\mathcal{F}_t = \mathcal{F}''$  in  $H_t = G''$ , is a shortest contractible cycle in  $G$ .

We next show that the running time of the algorithm is  $O(E^2 \log E)$ . Finding the shortest non-separating cycle<sup>1</sup> takes  $O(E^2 \log E)$  because of Lemma 1(a). Finding the arcs  $\alpha_1, \dots, \alpha_k$  used to cut  $G'$  takes  $O(n \log n)$  because we apply Lemma 1(b) to  $G'$ , which has  $O(E)$  edges. We hence obtain a plane graph  $G''$  with  $O(E)$  edges. Finally, finding the shortest cycle in  $G''$  in compliance with  $\mathcal{F}''$  is can be done in  $O(E^2)$  time because of Corollary 4.  $\square$

## 5 Hardness of related problems

Our NP-hardness results hold for unweighted graphs, so we will restrict our discussion in this section to unweighted graphs. All reductions are from the problem INDEPENDENTSET, which is NP-hard [15]:

<sup>1</sup>For orientable surfaces, we could use the tight closed walk of [3], which can be found faster. However, this would reduce the final running time only to  $O(E^2)$ .




Figure 6: Reduction in Theorem 8.

Problem: INDEPENDENTSET.

Input: a graph  $H$ .

Task: find a largest independent set in  $H$ .

Consider an input graph  $H$  for INDEPENDENTSET, which we assume for simplicity has  $V(H) = \{1, 2, \dots, n\}$ . Our reductions use an auxiliary graph  $G_H$  embedded in the sphere, that depends on  $H$ . The construction of  $G_H$  starts with an empty graph, and proceeds as follows:

1. For each vertex  $v \in V(H)$  with neighbors  $u_1 < \dots < u_{deg(v)}$ , create a path  $P_v$  with  $deg(v) + 1$  new vertices, denoted  $v, v(u_1), v(u_2), \dots, v(u_{deg(v)})$ , and with edges between  $v$  and  $v(u_1)$  and between  $v(u_i)$  and  $v(u_{i+1})$  for  $i = 1, \dots, deg(v) - 1$ . Add the path  $P_v$  to  $G_H$ .
2. For each vertex  $v \in V(H)$ , make a path  $Q_v$  with  $deg(v) + 2$  vertices, whose second vertex is denoted  $v'$ . Add the path  $Q_v$  to  $G_H$ .
3. For each vertex  $v \in V(H)$ , add new vertices  $s_v, t_v$  to  $G_H$ , add edges from  $s_v$  to the first vertex in  $P_v$  and the first vertex in  $Q_v$ , and add edges from  $t_v$  to the other ends of  $P_v$  and  $Q_v$ . This defines a cycle for each vertex  $v \in V(H)$ ; see Figure 6, left.
4. Add a new vertex  $s$  to  $G_H$ , add the edges  $ss_1, st_n$ , and add the edges  $t_i s_{i+1}$  for  $i = 1, \dots, n - 1$ ; see Figure 6, right.

This finishes the construction of  $G_H$ . The embedding of  $G_H$  in the sphere is fixed to be the one shown in Figure 6, right. In this embedding, the paths adjacent to  $s_i$  are ordered clockwise as  $s_i t_{i-1}, s_i Q_i, s_i P_i$  and the paths adjacent to  $t_i$  are ordered clockwise as  $t_i P_i, t_i Q_i, t_i s_{i+1}$ , where we use  $s = t_0 = s_{n+1}$ .

In Figure 7 there is an example showing  $G_H$  when  $H$  is  $K_4$  minus an edge.

Consider the family  $\mathcal{F}_{G_H}$  of forbidden pairs in  $G_H$  given by

$$\mathcal{F}_{G_H} = \{\{u(v), v(u)\} \mid uv \in E(H)\}.$$

This family  $\mathcal{F}_{G_H}$  of forbidden pairs has the following property: a cycle in  $G_H$  is in compliance with  $\mathcal{F}_{G_H}$  if and only if does not use both  $P_v$  and  $P_u$  for any  $uv \in E(H)$ . Let  $\mathcal{Q}_{G_H}$  be the set of cycles from  $G_H$  that are in compliance with  $\mathcal{F}_{G_H}$  and are of the form  $ss_1 R_1 t_1 s_2 R_2 t_2 \dots s_n R_n t_n s$ , where each  $R_v$  is either  $P_v$  or  $Q_v$ . The connection between  $H$  and  $G_H$  is summarized in the following observation.

**Lemma 7.** *The graph  $H$  has an independent set of size  $k$  if and only if the shortest cycle in  $\mathcal{Q}_{G_H}$  has length at most  $1 + 4n + 2 \cdot |E(H)| - k$ .*




Figure 7: Example of the reduction in Theorem 8. Left: the original graph  $H$ . Right: the graph  $G_H$ . For problem (a) in Theorem 8, the reduction asks for a cycle through  $s$  that is in compliance with  $\mathcal{F}_{G_H} = \{\{1(2), 2(1)\}, \{2(3), 3(2)\}, \{3(4), 4(3)\}, \{4(1), 1(4)\}, \{1(3), 3(1)\}\}$ . For problem (b) in Theorem 8, the reduction asks for a cycle in compliance with  $\mathcal{F}_{G_H} \cup \{\{1, 1'\}, \{2, 2'\}, \{3, 3'\}, \{4, 4'\}\}$ .

*Proof.* Assume first that  $H$  has an independent set  $I \subseteq \{1, \dots, n\}$  with  $k$  vertices. For  $v = 1, \dots, n$  let  $R_v$  be the path  $P_v$  if  $v \in I$  and  $Q_v$  if  $v \notin I$ . The cycle  $ss_1R_1t_1s_2R_2t_2 \cdots s_nR_nt_ns$  is in compliance with  $\mathcal{F}_{G_H}$  because there are no edges in  $H$  between the vertices  $I$ . Therefore this cycle is an element of  $\mathcal{Q}_{G_H}$  whose length is

$$\begin{aligned} 1 + \sum_{v=1}^n (2 + \# \text{vertices in } R_v) &= 1 + \sum_{v \in I} (2 + \# \text{vertices in } P_v) + \sum_{v \notin I} (2 + \# \text{vertices in } Q_v) \\ &= 1 + \sum_{v \in I} (2 + 1 + \deg(v)) + \sum_{v \notin I} (2 + 2 + \deg(v)) \\ &= 1 + 3|I| + 4(n - |I|) + \sum_{v=1}^n \deg(v) \\ &= 1 + 4n - k + 2 \cdot |E(H)|. \end{aligned}$$

To prove the other implication, assume that  $G_H$  has a cycle  $C \in \mathcal{Q}_{G_H}$  of length at most  $1 + 4n + 2 \cdot |E(H)| - k$ . Since  $C \in \mathcal{Q}_{G_H}$ , then  $C$  is of the form  $ss_1R_1t_1s_2R_2t_2 \cdots s_nR_nt_ns$ , where each  $R_v$  is either  $P_v$  or  $Q_v$ . Since  $C$  is in compliance with  $\mathcal{F}_{G_H}$ , the paths  $P_u$  and  $P_v$  cannot be in  $C$  if  $uv \in E(H)$ . Hence, the set  $I = \{v \in V(H) \mid R_v = P_v\}$  is an independent set of  $H$ . Using the bound on the length of  $C$  we then have

$$\begin{aligned} 1 + 4n + 2 \cdot |E(H)| - k &\geq (\text{length of } C) \\ &= 1 + \sum_{v \in I} (2 + \# \text{vertices in } P_v) + \sum_{v \notin I} (2 + \# \text{vertices in } Q_v) \\ &= 1 + 3|I| + 4(n - |I|) + \sum_{v=1}^n \deg(v) \\ &= 1 + 4n - |I| + 2 \cdot |E(H)|, \end{aligned}$$

and therefore  $|I| \geq k$ . □

As a consequence of the previous Lemma, searching for a largest independent set in  $H$  is equivalent to searching for a shortest cycle in  $\mathcal{Q}_{G_H}$ .

**Theorem 8.** *The following two problems are NP-hard for unweighted graphs:*

- (a) *Given a plane graph  $G$ , a family of forbidden pairs  $\mathcal{F}$ , which may be cofacial, and a vertex  $s$  of  $G$ , find a shortest cycle in compliance with  $\mathcal{F}$  that passes through  $s$ .*

- (b) Given a plane graph and a family of forbidden pairs  $\mathcal{F}$  that can be covered by 2 or more faces, find a shortest cycle in compliance with  $\mathcal{F}$ .

*Proof.* To show the NP-hardness of problem (a), consider the plane graph  $G_H$ , the family  $\mathcal{F}_{G_H}$  of forbidden pairs, and take  $s \in V(G_H)$  as the given vertex. The graph  $G_H$  is constructed such that any cycle through  $s$  is of the form  $ss_1R_1t_1s_2R_2t_2 \cdots s_nR_nt_ns$ , where each  $R_v$  is either  $P_v$  or  $Q_v$ . Hence, the family of cycles  $\mathcal{Q}_{G_H}$  is precisely the family of cycles in  $G_H$  through  $s$  that are in compliance with  $\mathcal{F}_{G_H}$ . Finding the shortest cycle in  $\mathcal{Q}_{G_H}$  would also solve the problem INDEPENDENTSET because of Lemma 7. This shows that problem (a) is NP-hard.

To show the NP-hardness of problem (b), consider the plane graph  $G_H$ , and the family  $\mathcal{F}'$  of forbidden pairs given by

$$\mathcal{F}' = \mathcal{F}_{G_H} \cup \{\{1, 1'\}, \{2, 2'\}, \dots, \{n, n'\}\}.$$

Note that the vertices in  $\mathcal{F}'$  are covered by two faces of  $G_H$ . Because of the ‘new’ forbidden pairs, a cycle in  $G_H$  in compliance with  $\mathcal{F}'$  cannot use both  $P_v$  and  $Q_v$  for any  $v \in V(H)$ . Inspection shows that, if  $P_v, Q_v$  cannot be used simultaneously, the only cycles in  $G_H$  are those of the form  $ss_1R_1t_1s_2R_2t_2 \cdots s_nR_nt_ns$ , where each  $R_v$  is either  $P_v$  or  $Q_v$ . It follows that the family of cycles in  $G_H$  in compliance with  $\mathcal{F}'$  is precisely  $\mathcal{Q}_{G_H}$ , and hence problem (b) is NP-hard.  $\square$

For studying contractible and separating cycles, the following observation will be useful.

**Lemma 9.** *Let  $G$  be a graph embedded in a surface  $\Sigma$  and let  $\alpha$  be a closed curve in  $\Sigma$  through a vertex  $v$  of  $G$ . If  $\alpha$  does not touch any other edge or vertex of  $G$  besides  $v$ , a separating cycle in  $G$  cannot cross  $\alpha$ .*

*Proof.* A separating cycle crosses each closed curve in  $\Sigma$  an even number of times. Since a cycle in  $G$  can have at most one point in common with  $\alpha$  by hypothesis, a separating cycle cannot cross  $\alpha$ .  $\square$

**Theorem 10.** *The following problem is NP-hard for unweighted graphs: given a graph  $G$  embedded in a surface and a vertex  $s \in V(G)$ , find a shortest contractible cycle that passes through  $s$ .*

*Proof.* Consider the graph  $G_H$  embedded in the sphere, as defined above. Let  $f_0$  be the face bounded by the cycle  $ss_1P_1t_1s_2P_2t_2 \cdots s_nP_nt_ns$ . We convert  $G_H$  into a graph embedded in an orientable surface of genus  $|E(H)|$ , as follows. For each vertex  $v(u)$  of  $V(G_H)$  take a small simple closed curve  $\alpha_{v(u)}$  passing through the vertex  $v(u)$  and that is otherwise contained in the interior of  $f_0$ . Moreover, the curves  $\alpha_{v(u)}$  should be pairwise disjoint for all  $v(u) \in V(G_H)$ . Let  $\tilde{\Sigma}$  be the surface obtained by removing the interior of the two disks bounded by  $\alpha_{v(u)}, \alpha_{u(v)}$ , for all  $uv \in E(H)$ . Then, for each edge  $uv \in E(H)$  we glue the two boundaries defined by  $\alpha_{v(u)}$  and  $\alpha_{u(v)}$ , in such a way that the vertices  $u(v), v(u)$  are identified and the resulting surface is orientable; see Figure 8. This finishes the transformation of  $G_H$ . Let  $G'_H$  be the resulting embedded graph, which is embedded in a certain surface  $\Sigma'$ . Since we add  $|E(H)|$  ‘independent handles’, the surface  $\Sigma'$  has genus  $|E(H)|$  and no boundary.

First, note that a separating cycle in  $G'_H$  cannot cross  $\alpha_{u(v)} = \alpha_{v(u)}$  because of Lemma 9 applied to the curve  $\alpha_{u(v)}$  in the surface  $\Sigma'$ . Since cutting the surface  $\Sigma'$  along the curves  $\alpha_{u(v)}$ , for all  $uv \in E(H)$ , give rise to  $\tilde{\Sigma}$ , it follows that a cycle in  $G'_H$  that is separating must be a cycle in  $G_H$  that is in compliance with  $\mathcal{F}_{G_H}$ . As a particular case, any contractible cycle in  $G'_H$  must be a cycle in  $G_H$  that is in compliance with  $\mathcal{F}_{G_H}$ . Note also that any




Figure 8: Illustration for the local transformation in the proof of Theorem 10. Left: part of the original graph  $G_H$  and two curves  $\alpha_{v(u)}$  and  $\alpha_{u(v)}$ . Right: part of the resulting surface  $\Sigma$  after adding the handle corresponding to the edge  $uv$  of  $H$ .

cycle in  $G_H$  becomes a contractible cycle in  $G'_H$  because  $f_0$  is the only face affected by the topological surgery.

It follows that finding a shortest contractible cycle in  $G'_H$  through  $s$  is equivalent to finding a shortest cycle in  $G_H$  through  $s$  that is in compliance with  $\mathcal{F}_{G_H}$ . As seen in the proof of Theorem 8(a), this is equivalent to finding a shortest cycles in  $\mathcal{Q}_{G_H}$ , which in turn is equivalent to solving the problem INDEPENDENTSET in the graph  $H$  because of Lemma 7. This shows that the problem is NP-hard. In general, the embedded graph we have constructed is not a cellular embedding. However, we can transform it to a cellular embedding by adding appropriate edges and subdividing each of them  $n^2$  times, so that they do not participate in any shortest contractible cycle.  $\square$

**Theorem 11.** *The following problem is NP-hard for unweighted graphs: given a graph embedded in a surface, find a shortest surface-separating cycle.*

*Proof.* Consider the embedded graph  $G'_H$  used in the proof of Theorem 10. We further modify  $G'_H$ , as follows. For each  $u \in \{1, \dots, n\}$ , let  $f_u$  be the face bounded by the paths  $s_u P_u t_u$  and  $s_u Q_u t_u$ . Take a small simple curve  $\alpha_u$  (resp.  $\alpha'_u$ ) passing through vertex  $u$  (resp.  $u'$ ) of  $G'_H$  and that is otherwise contained in the interior of the face  $f_u$ . Then we apply a similar surgery as in the proof of Theorem 10: for each  $u \in \{1, \dots, n\}$  we remove the interior of the two disks bounded by  $\alpha_u, \alpha'_u$  and glue the resulting boundaries, defined by  $\alpha_u$  and  $\alpha'_u$ , in such a way that the vertices  $u, u'$  are identified and the resulting surface is orientable. Let  $G''_H$  be the resulting embedded graph. It has genus  $|E(H)| + n$  because we have added  $n$  ‘independent handles’ to  $G'_H$ .

The same argument that was used with  $\alpha_{u(v)}$  in the proof of Theorem 10 can be used with the curves  $\alpha_{u'}$ . Therefore, a separating cycle in  $G''_H$  corresponds to a cycle in  $G_H$  that is in compliance with

$$\mathcal{F}' = \mathcal{F}_{G_H} \cup \{\{1, 1'\}, \{2, 2'\}, \dots, \{n, n'\}\},$$

and, as discussed in the proof of Theorem 8(b), corresponds to a cycle in  $\mathcal{Q}_{G_H}$ . Moreover, any cycle in  $\mathcal{Q}_{G_H}$  corresponds to a separating in  $G''_H$  because any handle is attached to only one face of  $G_H$ . It follows that finding a shortest separating cycle in  $G''_H$  corresponds to finding a shortest cycle in  $\mathcal{Q}_{G_H}$ , which in turn is equivalent to solving the problem INDEPENDENTSET in the graph  $H$  because of Lemma 7. This shows that the problem is NP-hard. Again, we can ensure we have an embedded graph with a cellular embedding by adding appropriate edges and subdividing each of them  $n^2$  times.  $\square$

## 6 Conclusions

We have given an algorithm that solves the problem of finding the shortest cycle in a  $n$ -vertex plane graph with cofacial forbidden pairs in  $O(n^2 \log n)$  time. It may seem a priori that an improvement reducing the running time to  $O(n \log n)$  should not be difficult. However, when the set of forbidden pairs is empty the problem is that of finding a shortest cycle in a planar graph, which is equivalent to finding a min-cut in the dual multigraph. The currently best deterministic algorithm to solve the min-cut problem in planar multigraphs takes  $O(n \log^2 n)$  time [5] (c.f. [10]). Therefore, finding an algorithm to find in  $O(n \log n)$  time the shortest cycle in a plane graph with cofacial forbidden pairs would have further implications for other problems.

We have also shown that a shortest contractible cycle in an embedded graph can be found in polynomial time, while several similar-looking problems are NP-hard. It seems interesting to explore the difference between finding shortest cycles and shortest walks with topological properties.

## References

- [1] S. Cabello. Many distances in planar graphs. In *SODA '06: Proc. 17th Symp. Discrete Algorithms*, pages 1213–1220, 2006.
- [2] S. Cabello and E. W. Chambers. Multiple source shortest paths in a genus  $g$  graph. In *SODA '07: Proc. 18th Symp. Discrete Algorithms*, pages 89–97, 2007.
- [3] S. Cabello, M. DeVos, J. Erickson, and B. Mohar. Finding one tight cycle. Manuscript available at <http://www.imfm.si/preprinti/PDF/01047.pdf>; preliminary version in *SODA '08: Proc. 19th Symp. Discrete Algorithms*.
- [4] S. Cabello and B. Mohar. Finding shortest non-separating and non-contractible cycles for topologically embedded graphs. *DCG*, 37:213–235, 2007. Preliminary version in *ESA '05*.
- [5] P. Chalermsook, J. Fakcharoenphol, and D. Nanongkai. A deterministic near-linear time algorithm for finding minimum cuts in planar graphs. In *SODA '04: Proc. 15th Symp. Discrete Algorithms*, pages 828–829, 2004.
- [6] E. W. Chambers, É. Colin de Verdière, J. Erickson, F. Lazarus, and K. Whittlesey. Splitting (complicated) surfaces is hard. *Computational Geometry: Theory & Applications*, 41:94–110, 2008. Preliminary version in *SOCG '06*.
- [7] É. Colin de Verdière and J. Erickson. Tightening non-simple paths and cycles on surfaces. In *SODA '06: Proc. 17th Symp. Discrete Algorithms*, pages 192–201, 2006.
- [8] É. Colin de Verdière and F. Lazarus. Optimal system of loops on an orientable surface. *DCG*, 33:507–534, 2005. Preliminary version in *FOCS '02*.
- [9] É. Colin De Verdière and F. Lazarus. Optimal pants decompositions and shortest homotopic cycles on an orientable surface. *J. ACM*, 54(4):18, 2007. Preliminary version in *Graph Drawing 2004*.
- [10] H. Djidjev. Computing the girth of a planar graph. In *ICALP*, volume 1853 of *LNCS*, pages 821–831. Springer, 2000.

- [11] D. Eppstein. Dynamic generators of topologically embedded graphs. In *SODA '03: Proc. 14th Symp. Discrete Algorithms*, pages 599–608, 2003.
- [12] J. Erickson and S. Har-Peled. Optimally cutting a surface into a disk. *DCG*, 31:37–59, 2004. Preliminary version in SOCG '02.
- [13] J. Erickson and K. Whittlesey. Greedy optimal homotopy and homology generators. In *SODA '03: Proc. 16th Symp. Discrete Algorithms*, pages 1038–1046, 2005.
- [14] H. N. Gabow, S. N. Maheswari, and L. J. Osterweil. On two problems in the generation of program test paths. *IEEE Trans. Software Eng.*, 2(3):227–231, 1976.
- [15] M. R. Garey and D. S. Johnson. *Computers and Intractability: A Guide to the Theory of NP-Completeness*. W.H. Freeman and Company, 1979.
- [16] A. Hatcher. *Algebraic Topology*. Cambridge University Press, 2001. Available at <http://www.math.cornell.edu/~hatcher/>.
- [17] M. Henzinger, P. Klein, S. Rao, and S. Subramanian. Faster shortest-path algorithms for planar graphs. *Journal of Computer and System Sciences*, 55(1):3–23, 1997.
- [18] V. Kann. Polynomially bounded minimization problems that are hard to approximate. *Nordic J. Comp.*, 1:317–331, 1994. Preliminary version in ICALP '93.
- [19] M. Kaufmann and D. Wagner, editors. *Drawing Graphs, Methods and Models*, volume 2025 of *Lecture Notes in Computer Science*. Springer, 2001.
- [20] M. Kutz. Computing shortest non-trivial cycles on orientable surfaces of bounded genus in almost linear time. In *SOCG '06: Proc. 22nd Symp. Comput. Geom.*, pages 430–438, 2006.
- [21] W. S. Massey. *Algebraic Topology: An Introduction*. Springer Verlag, 1967.
- [22] B. Mohar and C. Thomassen. *Graphs on Surfaces*. Johns Hopkins University Press, Baltimore, 2001.
- [23] T. Nishizeki and M. S. Rahman. *Planar Graph Drawing*, volume 12 of *Lecture Notes Series on Computing*. World Scientific Publishing Company, September 2004.
- [24] H. Ren and M. Deng. Short cycle structures for graphs on surfaces and an open problem of Mohar and Thomassen. *Science in China Series A: Mathematics*, 49(2):212–224, 2006.
- [25] J. Stillwell. *Classical Topology and Combinatorial Group Theory*. Springer-Verlag, New York, 1993.
- [26] C. Thomassen. Embeddings of graphs with no short noncontractible cycles. *J. Combin. Theory, Ser. B*, 48:155–177, 1990.
- [27] C. Thomassen. Recent results on graph embeddings. In Y. Alavi, G. Chartrand, O. R. Ollerman, and A. J. Schwenk, editors, *Graph Theory, Combinatorics, and Applications*, pages 1093–1103. John Wiley and Sons, 1991.