
Parabolic double cosets in Coxeter groups

Sara Billey^{*1}, Matjaz Konvalinka^{*2}, T. Kyle Petersen^{*3}, William Slofstra^{*4}, and Bridget Tenner^{*5}

¹Department of Mathematics, University of Washington, USA – United States

²Department of Mathematics, University of Ljubljana Institute for Mathematics, Physics and Mechanics – Ljubljana, Slovenia

³Dept of Mathematical Sciences, DePaul University – Chicago, IL 60614, United States

⁴Institute for Quantum Computing, Quantum-Nano Centre, University of Waterloo – Canada

⁵Dept of Mathematical Sciences, DePaul University – United States

Abstract

Parabolic subgroups WI of Coxeter systems (W, S) and their ordinary and double cosets W/WI and WI/WJ appear in many contexts in combinatorics and Lie theory, including the geometry and topology of generalized flag varieties and the symmetry groups of regular polytopes. The set of ordinary cosets wWI , for $I \subseteq S$, forms the Coxeter complex of W , and is well-studied. In this extended abstract, we look at a less studied object: the set of all double cosets $WIwWJ$ for $I, J \subseteq S$. Each double coset can be presented by many different triples (I, w, J) . We describe what we call the lex-minimal presentation and prove that there exists a unique such choice for each double coset. Lex-minimal presentations can be enumerated via a finite automaton depending on the Coxeter graph for (W, S) . In particular, we present a formula for the number of parabolic double cosets with a fixed minimal element when W is the symmetric group S_n . In that case, parabolic subgroups are also known as Young subgroups. Our formula is almost always linear time computable in n , and the formula can be generalized to any Coxeter group.

*Speaker